arXiv:1506.02412v1 [math.DS] 8 Jun 2015

ON THE ASYMPTOTIC WAVENUMBER OF SPIRAL WAVES IN $\lambda-\omega$ SYSTEMS.

M. AGUARELES, I. BALDOMÀ & T. M-SEARA

Abstract. In this paper we consider spiral wave solutions of a general class of $\lambda-\omega$ systems with a small parameter q and we prove that the asymptotic wavenumber of the spirals is a $\mathcal{C}^\infty$-flat function of the perturbation parameter q.

1. Introduction

Rigidly rotating spiral waves are commonly found in many chemical systems and biological processes [Kur84, LJD91, Win72, ZZ70]. In particular they are most likely to occur in oscillatory models having a rotational symmetry, such as generic $\lambda-\omega$ systems [Kur84], [Sch98]. These can be derived as the normal form of oscillatory reaction-diffusion systems near a Hopf bifurcation and read:

$$u_t = \Delta u + \lambda(f)u - \omega(f)w, \tag{1}$$

$$w_t = \Delta w + \omega(f)u + \lambda(f)w, \tag{2}$$

where $u = u(x, y, t)$, $v = v(x, y, t)$ and Δ denotes the Laplacian. λ and ω are real functions of $f = \sqrt{u^2 + w^2}$. The conditions that λ usually satisfies are: $\lambda(1) = 0$, to ensure that the system has a space independent limit solution and $\lambda' < 0$, to guarantee that this limit cycle is stable to homogeneous (space independent) perturbations. As for ω, based on stability considerations (see [NK81]), it is usually assumed that $|\omega'|$ is small.

Numerical computations reveal that the system (1)-(2) exhibits solutions in the shape of n-spirals (see for instance [BHO97, GB05]) and more precisely, in the shape of Archimedian spiral waves with a specific frequency Ω. These rigidly rotating solutions of (1)-(2) can then be written like

$$\begin{aligned} u(r,\phi,t) &= f(r)\cos\left(\Omega t + n\phi - \int_0^r v(s)\,ds\right), \\ w(r,\phi,t) &= f(r)\sin\left(\Omega t + n\phi - \int_0^r v(s)\,ds\right), \end{aligned} \tag{3}$$

being r and ϕ the polar radius and azymuthal coordinates of the plane and thus the Laplacian can be expressed as $\Delta = \partial_{rr} + \partial_r/r + \partial_{\phi\phi}/r^2$. Therefore, since $f(r)$ plays the role of a modulus, $f(r) \geq 0$ $\forall r > 0$ and also $f(0) = 0$ in order for u and w to be regular at $r = 0$. Also, in

---

Date: June 9, 2015.

order for these functions to have the shape of a spiral, the phase must increase or decrease monotonically as one moves away from the centre of the spiral and so $v(r)$, which is usually denoted as the *local wavenumber*, must have a constant sign for all r. In the particular case where $n = 0$, the phase is purely radial and they are usually denoted as *target patterns* since the lines of constant phase become concentric rings, that is to say, along any radial line, the pattern is asymptotically that of a plane wave.

Substituting the particular expressions (3) in (1)-(2) one obtains a set of ordinary differential equations in terms of the radial polar variable, r, that reads

$$0 = f'' + \frac{f'}{r} - f\frac{n^2}{r^2} + f(\lambda(f) - v^2), \tag{4}$$

$$0 = fv' + \frac{fv}{r} + 2f'v + f(\Omega - \omega(f)). \tag{5}$$

We note that any arbitrary constant can be added to the phase of the sine and cosine functions of u and w in (3) and they would still yield the same equations (4)-(5).

Using the identity

$$fv' + \frac{fv}{r} + 2f'v = \frac{(f^2vr)'}{rf}, \tag{6}$$

along with the fact that $(f^2(r)v(r)r)_{|r=0} = 0$, equation (5) can be expressed in the integral form,

$$v(r) = (rf^2(r))^{-1}\int_0^r tf^2(t)(\omega(f(t)) - \Omega)\, dt, \tag{7}$$

and this yields $v(0) = 0$.

Archimedian spiral waves are characterized by the fact that the distance between two neighbouring fronts of the isophase lines tends to a constant, as $r \to \infty$. That is to say, if we consider two points of an isophase line ($n\phi - \int_0^r v(s)\, ds = ctant$) one with coordinates (ϕ, r) and the following one on the same radial line with coordinates $(\phi + 2\pi, r + \delta(r))$, one obtains,

$$n\phi + \int_0^r v(s)\, ds = n(\phi + 2\pi) + \int_0^{r+\delta(r)} v(s)\, ds.$$

The separation between these two fronts is thus here represented by $\delta(r)$ and satisfies

$$\int_r^{r+\delta(r)} v(s)\, ds = 2\pi n.$$

Then, for Archimedian spiral waves it is expected that $\delta(r) \to D < \infty$ as $r \to \infty$. Using the mean value theorem in the last equality gives $v(r) \to v_\infty < \infty$ as $r \to \infty$ with $v_\infty = 2\pi n/D$, that is to say, v_∞ is proportional to the inverse of the spirals' front separation D, and it is usually known as the *asymptotic wavenumber*. As for the modulus, $f(r)$, the type of solutions that have been observed are such that $f(r)$ has a bounded limit and $f'(r) \to 0$ as $r \to \infty$. We will therefore focus on solutions of (4)-(5) such that $f(0) = 0$, $f(r)$ and $v(r)$ have bounded limits and $f'(r) \to 0$ as $r \to \infty$. These are four restrictions to a third order system of differential equations which suggests that there exists a selection mechanism for the frequency Ω, that is to say, Ω cannot be arbitrary. In fact, Kopell & Howard in [NK81] establish the existence of spiral wave solutions ($n \neq 0$) and target patterns ($n = 0$) for a particular value of Ω when $\lambda(1) = 0$, $\lambda'(x) < 0$ and $\omega(x) < 0$. These are then solutions of the system (4)-(5) for $n \in \mathbb{Z}$ with precisely the boundary conditions described above, that is,

$$\begin{aligned} &f(0) = v(0) = 0, \\ &\lim_{r\to\infty} f(r) = f_\infty < \infty, \quad \lim_{r\to\infty} f'(r) = 0, \quad \lim_{r\to\infty} v(r) = v_\infty < \infty \\ &f(r) > 0 \text{ and } v(r) \text{ has constant sign } \forall r > 0. \end{aligned} \tag{8}$$

Spiral wave solutions of systems of the type in (4)-(5) with some particular functions $\lambda(x)$, $\omega(x)$ have been studied by numerous researchers. For instance, Hagan in [Hag82] considers the particular case of the complex Ginzburg-Landau equation where $\lambda(x) = 1-x^2$ and $\omega(x) = -qx^2$. He uses the method of matching formal asymptotic expansions to construct spiral wave solutions for small values of the parameter q. In particular, he formally finds that the asymptotic wavenumber $v_\infty = \lim_{r\to\infty} v(r)$ and $\Omega - q$ are exponentially small in q. Also, Greenberg in [Gre81] uses a formal perturbation technique to construct solutions of (9)-(10) when $\lambda(x) = 1-x$ and $\omega(x) = 1 + q(x-1)$. Kopell and Howard in [NK81] establish the existence of spiral wave solutions of (4)-(5) under the hypothesis that $\lambda(1) = 0$, $\lambda'(\cdot) < 0$, $\omega'(\cdot) < 0$ and $|\omega'(\cdot)| << 1$.

The ultimate motivation of the work in this paper is precisely to investigate the exponentially small character of the asymptotic wavenumber for a general class of $\lambda - \omega$ systems. In particular, in this paper we consider a general class of $\lambda - \omega$ systems with the following conditions:

(A1) λ and ω belong to $\mathcal{C}^\infty(\mathbb{R})$ and they are such that $\lambda(1) = 0$, $\lambda'(1) < 0$, and $|\omega'| << 1$. We remark that by suitably rescaling the radius variable r and the phase function v a new function $\bar\lambda$ may be written such that $\bar\lambda(0)$ has any prescribed value. Therefore, and without lost of generality we also assume that $\lambda(0) = 1$.

(A2) $x\lambda(x)$ is concave, that is to say, $\partial_x^2(x\lambda(x)) < 0$.

Since we assume that $|\omega'| << 1$, we shall write $\omega(x) = \omega_0 + q\bar\omega(x)$, for $0 \leq q << 1$. We introduce a new parameter $\bar\Omega$ such that the frequency may also be written like $\Omega = \omega_0 + q\bar\Omega$. Dropping the bars to simplify the notation equations (4)-(5) read

$$0 = f'' + \frac{f'}{r} - f\frac{n^2}{r^2} + f(\lambda(f) - v^2), \tag{9}$$

$$0 = fv' + \frac{fv}{r} + 2f'v + qf(\Omega - \omega(f)), \tag{10}$$

where $0 < q << 1$ and $\Omega \in \mathbb{R}$ are the new parameters. As for the boundary conditions, we also consider the ones in [NK81] given in (8).

In this paper we prove that, for these solutions to exist, a necessary condition is that $\Omega = \Omega(q)$ has to be a $\mathcal{C}^\infty$ function of q such that $\partial_q^k\Omega(0) = 0$, for $k \geq 1$. To prove this result we provide a formal expansion in q of the solutions of equations (9)-(10) with boundary conditions (8). We obtain an infinite set of differential equations with suitable boundary conditions, one for each order in q, and we rigorously prove that all these equations have a unique bounded solution if and only if $\Omega(q) - \Omega(0)$ is a $\mathcal{C}^\infty$-flat function of q. Therefore, equations (9)-(10) with boundary conditions (8) can be solved up to any order in q. As a straightforward consequence of our results the solutions obtained in [NK81] satisfy that $\Omega - \Omega(0)$ and $v_\infty = \lim_{r\to\infty} v(r)$ are C^∞-flat functions of q. This is then a beyond all orders phenomenon and the rigorous study of the asymptotic values of Ω and v_∞ will be the goal of a forthcoming paper.

The paper is organised as follows. We start in Section §2 by posing the particular shape of the formal solution of (9)-(10). We then introduce our main theorem 2.2 of existence and uniqueness of this formal solution provided the power series for $|\Omega - \Omega(0)|$ has vanishing terms. We also provide a set of numerical computations for the classical complex Ginzburg-Landau problem where $\lambda(x) = 1 - x^2$ and $\omega(x) = -x^2$, which suggests that $\Omega - \Omega(0)$ is indeed not zero but exponentially small in q. In Section §3 we prove the main result as follows: we start by obtaining the equations and boundary conditions that each term in the asymptotic expansion satisfies and we then proceed by induction.

2. MAIN RESULT: A FORMAL SOLUTION

In this section we introduce and justify the expected particular form of the asymptotic expansion in q for the solution of system (9)-(10) with boundary conditions (8). We first start with a technical lemma that we shall prove in Section 2.2 below.

Lemma 2.1. *The system* (9)-(10) *with boundary conditions* (8) *has a solution* $(f(r), v(r))$ *if and only if*

$$v_\infty^2 - \lambda(f_\infty) = 0, \quad \omega(f_\infty) - \Omega = 0. \tag{11}$$

In addition, $\lim_{r\to\infty} v'(r) = 0$.

As a consequence, in order to have solutions of (9)-(10)-(8)*, the parameter* Ω *has to be a suitable function of* q*, i.e:* $\Omega = \Omega(q)$.

This theorem thus states that v_∞ and f_∞ are both functions of only the parameter q. To avoid cumbersome notation, we shall in general omit the dependence of q unless such omission leads to error.

Lemma 2.1 above implies that the solution of system (9)-(10) we are dealing with only depends on the small parameter q. We will call it $(f(r;q), v(r,q))$. By inspecting equations (9) and (10), one observes that the modulus $f(r;q)$, as well as the unknown frequency $\Omega(q)$, are even functions of q, that is $f(r;q) = f(r;-q)$ and $\Omega(q) = \Omega(-q)$, while v is an odd function of q, and so $v(r;q) = -v(r;-q)$. We can thus restrict to positive values of q without lost of generality. Moreover, using this even and odd character of the functions with respect to q we shall formally find the solutions to (9)-(10) as power series in q of the form:

$$\begin{aligned} f(r;q) &= \sum_{i\geq 0} f_i(r) q^{2i}, \quad v(r;q) = q\sum_{i\geq 0} v_i(r) q^{2i}, \\ \Omega(q) &= \sum_{i\geq 0} \Omega_i q^{2i}. \end{aligned} \tag{12}$$

In what follows we will find the differential equations and the boundary conditions that $f_k(r)$ and $v_k(r)$ have to satisfy. In order to solve these differential equations, we will find the terms for the expansion of the frequency, Ω_k.

Since we will deal with the behaviour as $r \to 0$ and $r \to +\infty$, we also introduce the notation

$$\psi = \mathrm{O}^l_m \Longleftrightarrow \psi(r) = \mathrm{O}\left(r^m\right), \ r \to 0 \text{ and } \psi(r) = \mathrm{O}\left(r^{-l}\right) \ r \to +\infty, \tag{13}$$

and

$$\psi = \mathrm{O}^{l,j}_m \Longleftrightarrow \psi(r) = \mathrm{O}\left(r^m\right), \ r \to 0 \text{ and } \psi(r) = \mathrm{O}\left(\log(r)^j r^{-l}\right) \ r \to +\infty, \tag{14}$$

which will be used along this paper without special mention.

The main result in this paper is:

Theorem 2.2. *Assume hypotheses (A1)-(A2) hold. Then the system* (9)-(10)-(8) *has a unique formal solution of the form* (12) *with* $\lim_{r\to+\infty} f_0(r) = 1$ *and satisfying that for all* $k \geq 0$*:*

$$f_k(0) = v_k(0) = 0 \text{ and } f_k(r), v_k(r) \text{ are bounded as } r \to \infty$$

if and only if

$$\Omega_0 = \omega(1), \quad \text{and} \quad \Omega_k = 0, \ \forall k \geq 1.$$

The functions $f_k(r), v_k(r)$ *also satisfy that*

$$\lim_{r\to\infty} v_0'(r) = 0, \quad \lim_{r\to\infty} v_0(r) = 0,$$

and

$$\lim_{r\to\infty} f_k(r) = \lim_{r\to\infty} v_k(r) = 0, \quad \text{for all } k > 0.$$

Moreover,

$$\begin{aligned} &f_0(r) = \mathrm{O}\left(r^n\right) \ \text{ as } r \to 0, \ 1 - f_0(r) = \mathrm{O}\left(r^{-2}\right) \ \text{ as } r \to +\infty. \\ &f_0' \in \mathrm{O}^3_{n-1}, \ f_0'' \in \mathrm{O}^4_{\max\{n-2,0\}} \\ &v_0 \in \mathrm{O}^{1,1}_1, \ v_0' \in \mathrm{O}^{2,1}_0, \ v_0'' \in \mathrm{O}^{3,1}_0 \end{aligned}$$

and for $k \geq 1$,

$$\begin{aligned} &f_k \in \mathrm{O}^{2,2k}_n, \ f_k' \in \mathrm{O}^{3,2k}_{n-1}, \ f_k'' \in \mathrm{O}^3_{\max\{n-2,0\}} \\ &v_k \in \mathrm{O}^{1,2k+1}_1, \ v_k' \in \mathrm{O}^{2,2k+1}_0, \ v_k'' \in \mathrm{O}^{3,2k+1}_0 . \end{aligned}$$

Finally if ω *is a monotone function,* v_0 *has constant sign.*

Remark 2.3. *Note that when* $q = 0$*, equation* (10) *becomes*

$$0 = fv' + \frac{fv}{r} + 2f'v = \frac{(f^2vr)'}{fr},$$

so $f^2(r;0)v(r;0)r \equiv c$*, which, upon evaluating at* $r = 0$ *gives* $c = 0$*. Since we are interested in non trivial solutions for* $f(r;0)$*, we obtain that* $v(r;0) \equiv 0$*. As a consequence,* $v_\infty = 0$*. Henceforth, equations* (11) *imply that* $\lambda(f_\infty) = 0$ *and* $\Omega = \omega(f_\infty)$*. Since by assumption (A1)* $\lambda(1) = 0$ *it seems natural to choose* $f_\infty = 1$*. In fact, if* $\lambda'(x) < 0$ *this is the only choice.*

Moreover, since $\lambda'(1) < 0$*, when* $q \neq 0$*, for any* $v_\infty = O(q)$*, the equation* $\lambda(f_\infty) = v_\infty^2$ *has a solution* $|f_\infty - 1| = O(q)$*, if* $|q|$ *is small enough.*

On the other hand, if $\lambda(x)$ *has another zero* $x_0 < 1$ *satisfying* $\lambda'(x_0) < 0$*, Theorem 2.2 can also be applied in this case by rescaling* $f = \overline{f}x_0$*. In conclusion, the condition* $\lim_{r\to+\infty} f_0(r) = 1$ *is not restrictive.*

From this theorem we conclude that, if system (9)-(10)-(8) has solution $(f(r;q), v(r;q))$ either $v_\infty = \lim_{r\to\infty} v(r) = 0$ or $v_\infty = \lim_{r\to\infty} v(r) = O(q^k)$, $\forall k \geq 0$. That is, the solution $v(r;q)$ either vanishes as $r \to \infty$ or it is C^∞-flat in q. In fact, numerical computations reveal that $v(r;q)$ is indeed not zero at infinity.

2.1. **Numeric computations.**

As an example we have considered a Ginzburg-Landau system for $n = 1$, which corresponds to $\lambda(x) = 1 - x^2$ and $\omega(x) = -x^2$, that is:

$$0 = f'' + \frac{f'}{r} - f\frac{1}{r^2} + f(1 - f^2 - v^2), \tag{15}$$

$$0 = fv' + \frac{fv}{r} + 2f'v + qf(\Omega - f^2). \tag{16}$$

with boundary conditions (8). We have used a MATLAB routine to obtain v_∞ which uses a finite difference scheme implementing the three-stage Lobatto Illa formula which provides a C^1-continuous solution that is fourth-order accurate uniformly in the interval of integration.

In [Hag82] and [ACW10] it is formally obtained an expression for v_∞ as a function of q, which is found to be of the form $v_\infty \sim Ae^{-B/q}q^{-1}$. Performing a linear fit of $\log(qv_\infty)$ with 95% confidence we obtain $B = 1.588191499224517$, using moderate values of $q \in [0.2, 0.5]$, so it seems to agree with the predicted value $B = \pi/2$. A rigorous computation of A and B would require working with multiprecision and it is beyond the scope of this paper.

2.2. **Dependence of Ω on the parameter q.**

We now prove Lemma 2.1. Assume that system $(9) - (10) - (8)$ has a solution $(f(r;q,\Omega), v(r;q,\Omega))$. We will omit the dependence on the parameters if there is no danger of confusion. To prove Lemma 2.1 we first recall that when $q = 0$, $v \equiv 0$, as it is pointed out in the Remark 2.3. As a consequence, $v_\infty = O(q)$.

Now we check that $f_\infty \neq 0$. Assume that $f_\infty = 0$. In this case, since $v_\infty = O(q)$, we have that $v_\infty << 1 = \lambda(0) = \lambda(f_\infty)$ taking q small enough. It follows that, for r_0 large enough



FIGURE 1. Value of $\log(qv_\infty)$ as a function of $1/q$, with $q \in [0.2, 0.5]$ for the system (15)-(16).

$n^2/r^2 < \lambda(f(r)) - v^2(r)$, for all $r \geq r_0$ and using that f is a positive solution of (9):

$$(rf'(r))' = rf''(r) + f'(r) < 0 \Longrightarrow f'(r) < \frac{r_0}{r} f'(r_0) \Longrightarrow f(r) - f(r_0) < r_0 f'(r_0) \log(r_0^{-1} r).$$

Since $f_\infty = 0$ and $f(r) > 0$, one can take r_0 large enough such that $f'(r) < 0$ for $r \geq r_0$. Taking $r \to \infty$ in the above inequality, we find a contradiction with the fact that f is a bounded function.

According to equation (9), using that $f'(r) \to 0$ as $r \to +\infty$, we obtain that $\lim_{r\to\infty} f''(r) = -f_\infty(\lambda(f_\infty) - v_\infty^2)$, while equation (10) provides that $\lim_{r\to\infty} v'(r) f_\infty = -q f_\infty(\Omega - \omega(f_\infty))$. Then, taking into account that $f_\infty \neq 0$ and the simple fact that:

$$\text{if } h(r) \text{ is bounded for all } r > 0 \text{ and } \lim_{r\to+\infty} h'(r) = b, \text{ then } b = 0, \tag{17}$$

which is immediate by Hôpital's rule, it is found that $\lim_{r\to+\infty} f''(r) = 0$ and $\lim_{r\to+\infty} v'(r) = 0$. Therefore, one is left with the couple of equations for the boundary values at infinity:

$$\lambda(f_\infty) - v_\infty^2 = 0, \quad \Omega - \omega(f_\infty) = 0.$$

Finally, if we explicitly write the dependence on q, Ω and we have that, to have solutions of our problem:

$$\chi(q, \Omega) := \Omega - \omega(f_\infty(q, \Omega)) = 0.$$

Note that, when $q = 0$, for any value of Ω, $v(r; 0, \Omega) = 0$ and f has to satisfy the equation

$$f'' + \frac{f'}{r} - f\frac{n^2}{r^2} + f\lambda(f) = 0$$

which is independent of Ω so that $f_\infty(0, \Omega) = f_\infty^0$ does not depend on Ω and thus $\partial_\Omega f_\infty(0, \Omega) = 0$. Therefore, using this and differentiating $\chi(q, \Omega)$ with respect to Ω one is left with $\partial_\Omega \chi(0, \Omega(f_\infty^0)) =$

$1 \neq 0$, which along with the fact that $\chi(0, \Omega(f_\infty^0)) = 0$, the implicit function theorem defines a function $\Omega(q)$ such that $\chi(q, \Omega(q)) = 0$ if $|q|$ is small enough.

3. POWER SERIES EXPANSIONS: PROOF OF THEOREM 2.2

The idea of the proof is as follows: we first start by describing the system of equations for f_k, v_k, $k \geq 0$ introduced in (12). We then deduce the boundary conditions that f_k, v_k must satisfy in order to be bounded solutions of such equations. We then prove the existence of f_k, v_k, along with some useful properties of the leading order terms f_0, v_0.

We emphasize that this is a formal procedure, so we do not pay special attention to some constants which, of course, could grow with respect to k at any formal step. For this reason we sometimes avoid the exact computation of some of these constants and we indeed may use the same letter to denote different ones.

3.1. Differential equations for f_k, v_k.

We are going to describe the equations that $\{f_k, v_k\}_{k \geq 0}$ have to satisfy. As it is usual, the equations for the leading order terms f_0 and v_0 will be nonlinear while the equations for f_k, v_k will be found to be non-homogeneous linear equations. To shorten the notation we introduce $F(x) = x\lambda(x)$ and $\tilde{\omega}(x) = x\omega(x)$ and we denote $DF(x)$ and $D\tilde{\omega}(x)$ to denote the derivatives respect to x of these functions.

With this notation, equations (9), (10) read:

$$0 = f'' + \frac{f'}{r} - f\frac{n^2}{r^2} + F(f) - fv^2, \tag{18}$$

$$0 = fv' + \frac{fv}{r} + 2f'v + q(f\Omega - \tilde{\omega}(f)), \tag{19}$$

and we consider the formal expansions defined in (12):

$$f(r;q) = \sum_{i \geq 0} f_i(r) q^{2i}, \quad v(r;q) = q \sum_{i \geq 0} v_i(r) q^{2i}, \quad \Omega(q) = \sum_{i \geq 0} \Omega_i q^{2i}.$$

Proposition 3.1. *The leading order terms f_0 and v_0, satisfy the equations*

$$0 = f_0'' + \frac{f_0'}{r} - n^2 \frac{f_0}{r^2} + F(f_0), \tag{20}$$

$$0 = f_0 v_0' + \frac{f_0 v_0}{r} + 2f_0' v_0 + f_0 \Omega_0 - \tilde{\omega}(f_0). \tag{21}$$

For $k \geq 1$, f_k and v_k satisfy the linear nonhomogeneous equations:

$$f_k'' + \frac{f_k'}{r} - n^2 \frac{f_k}{r^2} + DF(f_0) f_k = b_k(r), \tag{22}$$

$$f_0 v_k' + \frac{f_0 v_k}{r} + 2f_0' v_k + f_0 \Omega_k = c_k(r), \tag{23}$$

where

$$\begin{aligned} b_k(r) = &- \sum_{i=1}^{k} D^{2i} F(f_0(r)) \sum_{\substack{k_1 + \cdots + k_i = k \\ 1 \leq k_j \leq k-1}} f_{k_1}(r) \cdots f_{k_i}(r) \\ &+ \sum_{i=1}^{k} \sum_{l=1}^{i} f_{k-i}(r) \cdot v_{l-1}(r) \cdot v_{i-l}(r), \end{aligned} \tag{24}$$

$$\begin{aligned} c_k(r) = &\sum_{i=0}^{k-1} \big(f_{k-i}(r) \big(v_i'(r) + r^{-1} v_i(r) \big) + 2 f_{k-i}'(r) v_i(r) \big) + \sum_{i=0}^{k-1} f_{k-i}(r) \cdot \Omega_i \\ &- \sum_{i=1}^{2k} D^{2i} \tilde{\omega}(f_0(r)) \sum_{\substack{k_1 + \cdots + k_i = k \\ 1 \leq k_j}} f_{k_1}(r) \cdots f_{k_i}(r) \end{aligned} \tag{25}$$

with $F(x) = x\lambda(x)$ *and* $\tilde{\omega}(x) = x\omega(x)$*. In particular,* b_k *is independent of* f_k *and* v_k *and* c_k *is independent of* v_k*.*

Proof. By substituting expression (12) in (18), one obtains equation (20) for f_0. As for v, equation (19), gives to leading order equation (21) for v_0.

We now deal with $f_k, v_k,\ k \geq 1$. To illustrate the procedure we start by obtaining the particular equations for f_1, v_1. Expanding equation (18) in powers of q, the order $\mathrm{O}\left(q^2\right)$ provides an equation for f_1 in terms of v_0 and f_0, which reads,

$$f_1'' + \frac{f_1'}{r} - n^2 \frac{f_1}{r^2} + DF(f_0) f_1 = f_0 v_0^2, \tag{26}$$

which gives $b_1(r) = f_0(r) v_0(r)^2$.

Expanding equation (19) in powers of q, the order $\mathrm{O}\left(q^3\right)$ provides an equation for v_1 in terms of f_0, f_1 and v_0:

$$f_0 v_1' + \frac{f_0 v_1}{r} + 2 f_0' v_1 + f_0 \big(\Omega_1 - f_1 D\omega(f_0) \big) = c_1(r) \tag{27}$$

with

$$c_1(r) = -f_1(r)(\Omega_0 - \omega(f_0(r))) - f_1(r) v_0'(r) - r^{-1} f_1(r) v_0(r) - 2 v_0(r) f_1'(r). \tag{28}$$

To deal with the general case we first observe that the ansatz (12) may also be expressed in terms of a Taylor expansion of f, v and Ω with respect to q. Using the expansions (12), we deduce that $\partial_q^{2i+1} f(r;0) = 0$, $\partial_q^{2i} v(r;0) = 0$ and $\partial_q \Omega^{2i+1}(0) = 0$. This yields:

$$\begin{aligned} f(r;q) &= \sum_{i\geq 0} \frac{\partial_q^{2i} f(r;0)}{(2i)!} q^{2i}, \quad v(r;q) = q \sum_{i\geq 0} \frac{\partial_q^{2i+1} v(r;0)}{(2i+1)!} q^{2i}, \\ \Omega(q) &= \sum_{i\geq 0} \frac{\partial_q^{2i}\Omega(0)}{(2i)!} q^{2i}. \end{aligned} \tag{29}$$

Therefore,

$$f_k(r) = \frac{\partial_q^{2k} f(r;0)}{(2k)!}, \quad v_k(r) = \frac{\partial_q^{2k+1} v(r;0)}{(2k+1)!} \quad \text{and} \quad \Omega_k = \frac{\partial_q^{2k}\Omega(0)}{(2k)!}. \tag{30}$$

As a consequence, in order to obtain the equations for f_k, v_k and a general expression for b_k, c_k it is enough to differentiate equations (18) and (19) with respect to q. We shall use Faa di Bruno formula along with Leibnitz's rule.

We first deal with (18). We must compute the $2k$- derivative with respect to q of the nonlinear term $F(f) - fv^2$ and then evaluate at $q = 0$. Using Faa di Bruno's formula and the identity (30) gives

$$\begin{aligned} \frac{\partial_q^{2k}(F\circ f)(r;0)}{(2k)!} =& \sum_{i=1}^{2k} D^i F(f(r;0)) \sum_{\substack{k_1+\cdots+k_i = 2k \\ 1\leq k_j}} \frac{\partial_q^{k_1} f(r;0)}{k_1!} \cdots \frac{\partial_q^{k_i} f(r;0)}{k_i!} \\ =& DF(f_0(r)) f_k(r) + b_k^1(r), \end{aligned}$$

where, upon using once more identity (30) along with $\partial_q^{2l+1} f(r;0) = 0$, b_k^1 is found to read:

$$b_k^1(r) := \sum_{i=1}^{k} D^{2i} F(f_0(r))) \sum_{\substack{k_1+\cdots+k_i = k \\ 1\leq k_j \leq k-1}} f_{k_1}(r)\cdots f_{k_i}(r). \tag{31}$$

We note that the last sum does only depend on f_l with $0 < l < k$.

We now proceed likewise with $\partial_q^{2k}(f\cdot v^2)(r;0)$. Here we also note that $\partial_q^{2l} v(r;0) = 0$. Then, using Leibnitz rule:

$$\begin{aligned} b_k^2(r) :&= \partial_q^{2k}(f\cdot v^2)(r;0) = \sum_{i=2}^{2k} \begin{pmatrix} 2k \\ i \end{pmatrix} \partial_q^{2k-i} f(r;0)\cdot \partial_q^i v^2(r;0) \\ &= \sum_{j=1}^{k}\sum_{m=1}^{2j-1} \begin{pmatrix} 2k \\ 2j \end{pmatrix} \begin{pmatrix} 2j \\ m \end{pmatrix} \partial_q^{2k-2j} f(r;0)\cdot \partial_q^m v(r;0)\cdot \partial_q^{2j-m} v(r;0) \\ &= (2k)! \sum_{j=1}^{k}\sum_{l=1}^{j} f_{k-j}(r)\cdot v_{l-1}(r)\cdot v_{j-l}(r), \end{aligned} \tag{32}$$

so b_k^2 only depends on f_l, v_l with $0 \le l < k$.

Using the above expressions for $\partial_q^{2k}(F(f) - fv^2)$, we compute the $2k$-derivative of equation (18) with respect to q and, evaluating this derivative at $q = 0$, one finds that $f_k(r) = \partial_q^{2k} f(r;0)/(2k)!$ is a solution of the linear equation

$$f_k''(r) + \frac{f_k'(r)}{r} - f_k(r)\frac{n^2}{r^2} + DF(f_0(r)) \cdot f_k(r) + b_k^1(r) - \frac{b_k^2(r)}{(2k)!} = 0.$$

Therefore, f_k satisfies equation (22) with $b_k = -b_k^1 + b_k^2/(2k)!$ having the form (24).

We now deal with equation (19). The procedure is exactly analogous to the one for equation (18). First, we observe that, using the Leibnitz's rule, as well as identity (30),

$$\begin{aligned}\partial_q^{2k+1}&\left(f(r;0) \cdot v'(r;0) + \frac{f(r;0) \cdot v(r;0)}{r} + 2f'(r;0) \cdot v(r;0)\right)\\ &= (2k+1)!\big[f_0(r)\big(v_k'(r) + \frac{v_k(r)}{r}\big) + 2f_0'(r)v_k(r)\big] + c_k^1(r),\end{aligned}$$

where,

$$c_k^1(r) = (2k+1)!\sum_{i=0}^{k-1}\left(f_{k-i}(r)\big(v_i'(r) + \frac{v_i(r)}{r}\big) + 2f_{k-i}'(r)v_i(r)\right).$$

It only remains to compute the $2k+1$-derivative with respect to q of the nonlinear term $qf\big(\Omega(q) - \omega(f)\big)$. First, we define $\widetilde{\Omega}(q) = q\Omega(q)$ and we compute $\partial_q^{2k+1}\big(f(r;0) \cdot \widetilde{\Omega}(0)\big)$. We obtain, using Leibnitz's rule,

$$\begin{aligned}\partial_q^{2k+1}\big(f(r;0) \cdot \widetilde{\Omega}(0)\big) &= \sum_{i=0}^{2k+1}\left(\begin{array}{c} 2k+1 \\ i \end{array}\right)\partial_q^{2k+1-i} f(r;0) \cdot \partial_q^i \widetilde{\Omega}(0)\\ &= (2k+1)! f_0 \Omega_k + c_k^2(r),\end{aligned}$$

where

$$c_k^2(r) = (2k+1)!\sum_{i=0}^{k-1} f_{k-i}(r) \cdot \Omega_i.$$

We now introduce $\tilde{\omega}(x) = x\omega(x)$ and compute $c_k^3(r) := \partial_q^{2k+1}\big(q\tilde{\omega}(f(r;q))\big)_{|q=0}$:

$$\begin{aligned}
\frac{c_k^3(r)}{(2k+1)!} &= \frac{1}{(2k)!}\partial_q^{2k}\big(\tilde{\omega}(f(r;0))\big) \\
&= \sum_{i=1}^{2k} D^i\tilde{\omega}(f(r;0)) \sum_{\substack{k_1+\cdots+k_i=2k \\ 1\le k_j}} \frac{\partial_q^{k_1}f(r;0)}{k_1!}\cdots\frac{\partial_q^{k_i}f(r;0)}{k_i!} \\
&= \sum_{i=1}^{2k} D^i\tilde{\omega}(f_0(r)) \sum_{\substack{k_1+\cdots+k_i=k \\ 1\le k_j}} f_{k_1}(r)\cdots f_{k_i}(r)
\end{aligned}$$

Finally we compute the $2k+1$-derivative with respect to q of equation (19) and we obtain that $v_k(r) = \partial_q^{2k+1}v(r;0)/(2k+1)!$ satisfies equation (23) with c_k defined as in (25). □

3.2. **The leading order term.** We have already proved that the leading order terms f_0 and v_0, along with a suitable choice of $\Omega_0 = \omega(1)$ have to be solutions of the boundary problem:

$$\begin{aligned}
&0 = f_0'' + \frac{f_0'}{r} - n^2\frac{f_0}{r^2} + f_0\lambda(f_0) \\
&f_0(0) = 0, \quad \lim_{r\to\infty} f_0(r) = 1,
\end{aligned} \tag{33}$$

and

$$\begin{aligned}
&0 = f_0v_0' + \frac{f_0v_0}{r} + 2f_0'v_0 + f_0(\Omega_0 - \omega(f_0)) \\
&v_0(0) = 0, \quad \lim_{r\to\infty} v_0(r) < +\infty.
\end{aligned} \tag{34}$$

It is clear that the nonlinear equation for f_0 is qualitatively different to the ones for f_k with $k \ge 1$, which are all of them nonhomogeneous linear equations. Moreover, in order to begin an induction procedure (which will be our strategy to prove Theorem 2.2) we also need to prove the existence and properties of v_0. For that reason we study the leading order terms separately. Next proposition proves the part of Theorem 2.2 related to f_0 and v_0.

Proposition 3.2. *The boundary problem* (33) *has a bounded solution* $f_0 > 0$*. Moreover,* f_0 *satisfies the following inequalities*

$$0 < rf_0'(r) \le n^2 f_0(r), \qquad r > 0,$$

and it has the asymptotic expansions,

$$f_0(r) = \alpha r^n + \mathrm{O}\left(r^{n+1}\right), \ \text{ as } r \to 0 \ \text{ and } \ f_0(r) = 1 - \frac{n^2}{dr^2} + \mathrm{O}\left(r^{-4}\right), \ \text{ as } r \to +\infty. \tag{35}$$

with $d = -\lambda'(1)$*. We also have that* $f_0' \in \mathrm{O}_{n-1}^3$ *with* $\lim_{r\to+\infty} r^3 f_0'(r) = 2\frac{n^2}{d}$ *and* $f_0'' \in \mathrm{O}_{\min\{0,n-2\}}^4$.

The problem (34) *has a bounded solution* v_0 *if and only if* $\Omega_0 = \omega(1)$. *Moreover,*

$$v_0(r) = (r f_0^2(r))^{-1} \int_0^r t f_0(t)^2 (\omega(f_0(t)) - \Omega_0)\, dt, \tag{36}$$

and it satisfies the asymptotic expansions

$$v_0(r) = Cr + \mathrm{O}\left(r^2\right) \quad \text{as } r \to 0, \quad v_0(r) = -\frac{n^2 \omega'(1) \log(r)}{dr} + \mathrm{O}\left(1/r\right) \quad \text{as } r \to +\infty. \tag{37}$$

Moreover, $v_0' \in \mathrm{O}_0^{2,1}$, $v_0'' \in \mathrm{O}_0^{3,1}$.

When ω *is a monotone function the solution* v_0 *has constant sign.*

Proof. As it is shown in [AB11], the boundary problem (33) has a unique bounded solution. The inequalities and the expansions for $f_0(r)$ were rigourosly proven in [AB11] for the case that λ is an analytic function (Section 2 as $r \to 0$ and Sections 4 and 5 as $r \to +\infty$). The expansion for r small enough is also true in the case $\lambda \in \mathcal{C}^\infty$ and the behaviour as $r \to +\infty$ can be straightforwardly deduced from Lemma 2.14 and Remark 2.15 in [AB11]. Then, re-writting equation (33) we get the identity $(r f_0'(r))' = n^2 r^{-1} f_0(r) - f_0(r) \lambda(f_0(r))$. From this identity we deduce the asymptotic expansions for f_0' and f_0''.

We also know that $\lim_{r\to+\infty} r^3 f_0'(r)$ exists from the previous work in [AB11]. To compute this limit, we use L'Hôpital's rule, and the asymptotic expansion for f_0:

$$\frac{n^2}{d} = \lim_{r\to+\infty} r^2 \big(1 - f_0(r)\big) = \lim_{r\to+\infty} r^2 \int_r^{+\infty} f'(\xi)\, d\xi = \lim_{r\to+\infty} \frac{r^3 f'(r)}{2}$$

and so the results for f_0 are proven.

As for v_0, since $v_0(0) = 0$ and it satisfies equation (34), using property (6), gives to leading order expression (36). Now, using the asymptotic behaviour of $f_0(r)$ as $r \to \infty$ in (36), gives

$$\begin{aligned} v_0(r) =& \big(r - 2n^2/(d^2 r) + o(r^{-1})\big)^{-1} \left(\int_0^{r_0} t f_0(t)^2 (\omega(f_0(t)) - \Omega_0)\, dt \right. \\ &\left. + \int_{r_0}^r t \left(\omega(1) - \Omega_0 - \frac{n^2}{t^2\, d} \big(2(\omega(1) - \Omega_0) + \omega'(1)\big) + \mathrm{o}\left(t^{-2}\right) \right) dt \right), \end{aligned}$$

provided $r \geq r_0$ and r_0 is large enough. This last expression shows that in order for v_0 to be bounded at infinity, we have to impose $\omega(1) = \Omega_0$ and so this gives the asymptotic behaviour of $v_0(r)$ as $r \to \infty$ presented in (37).

Also, the asymptotic behaviour of $v_0(r)$ as $r \to 0$ is easily obtained by using the asymptotic expression of f_0 in equation (36),

$$v_0(r) \sim \frac{\int_0^r t \alpha^2 t^{2n} (\omega(\alpha t^n) - \omega(1))\, dt}{\alpha^2 r^{2n+1}} = \frac{\omega(0) - \omega(1)}{2n+2} r + \mathrm{O}\left(r^2\right).$$

The asymptotic behaviour of both v_0' and v_0'' follows from the fact that $v_0 \in \mathrm{O}_1^{1,1}$ is a solution of equation (34) along with the asymptotic behaviour of f_0, f_0'.

It only remains to check that $v_0(r)$ has constant sign when $\omega(x)$ is a monotone function. For instance, according to (36) if $\omega(x)$ is decreasing, since $\Omega_0 = \omega(1)$, $\omega(0) - \Omega_0 \geq \omega(f_0(t)) - \Omega_0 \geq \omega(1) - \Omega_0 = 0$, and hence $v_0(r) \geq 0$ for all $r \geq 0$. Likewise, if $\omega(x)$ is increasing, $v_0(r) \leq 0$ for all $r \geq 0$. $\square$

3.3. **Existence and properties of f_k. An induction procedure.** In this section we are going to prove the results of Theorem 2.2 related to f_k, v_k for $k \geq 1$. We will use the notation and results from Proposition 3.1. More precisely, we will prove that the problems:

$$\begin{aligned} &f_k'' + \frac{f_k'}{r} - n^2\frac{f_k}{r^2} + DF(f_0(r))f_k = b_k(r) \\ &f_k(0) = 0, \qquad f_k(r) \text{ bounded } r \geq 0 \end{aligned} \tag{38}$$

with $F(x) = x\lambda(x)$, and

$$\begin{aligned} &f_0 v_k' + \frac{f_0 v_k}{r} + 2f_0' v_k + f_0\Omega_k = c_k(r), \\ &v_k(0) = 0, \qquad v_k(r) \text{ bounded } r \geq 0 \end{aligned} \tag{39}$$

have solutions f_k and v_k provided

$$\Omega_k = 0. \tag{40}$$

Recall that b_k, c_k, were defined in Proposition 3.1, for $k \geq 1$. To prove this result we will use an induction procedure.

We first recall that, if $f_0, f_1 \cdots, f_{k-1}$ and $v_0, v_1, \cdots, v_{k-1}$ are known, then, the independent term b_k of (38) is determined and henceforth f_k satisfies a linear non-homogeneous equation. If we are able to prove the existence of such a solution, then, by property (6) and taking into account that $v_k(0) = 0$, we will have an explicit expression for v_k which depends on Ω_k and c_k:

$$v_k(r) = \big(rf_0^2(r)\big)^{-1}\int_0^r tf_0(t)\big(c_k(t) - f_0(t)\Omega_k\big)\,dt \tag{41}$$

Recall here that c_k depends only on $f_0, \cdots, f_k$ and $v_0, \cdots, v_{k-1}$.

Therefore, once one knows how to solve the equation for f_k, the function v_k is totally determined. Since all the equations for f_k have the same shape, it is mandatory to study the existence of solutions of linear equations of the form

$$\begin{aligned} &g''(r) + \frac{g'(r)}{r} - n^2\frac{g(r)}{r^2} + DF(f_0(r))g(r) = h(r) \\ &g(0) = 0, \qquad g(r) \text{ bounded } r \geq 0. \end{aligned} \tag{42}$$

We state the following technical lemma which will be proven in Subsection 3.4 by using the Fixed Point Theorem in a suitable Banach space.

Lemma 3.3. *Let $h : [0, +\infty) \to \mathbb{R}$ be a C^2 function. We define*

$$\mathcal{E}[h](r) := h''(r) + \frac{h'(r)}{r} - h(r)\frac{n^2}{r^2} + \big[DF(f_0(r)) + d\big]h(r)$$

with $F(x) = x\lambda(x)$ *and* $d = -\lambda'(1)$. *Assume that* $\mathcal{E}[h] \in \mathrm{O}^3_{n-1}$, *that is:*

$$\mathcal{E}[h](r) = \mathrm{O}\left(r^{n-1}\right), \quad r \to 0, \qquad \mathcal{E}[h](r)(r) = \mathrm{O}\left(r^{-3}\right), \quad r \to +\infty. \tag{43}$$

Then there exists a unique bounded solution g *of the boundary problem* (42). *Moreover, if* $\Delta g = g + hd^{-1}$, *we have that*

$$\Delta g \in \mathrm{O}^3_n, \quad \Delta g' \in \mathrm{O}^3_{n-1}, \quad \textit{and} \quad \Delta g'' \in \mathrm{O}^3_{\max\{n-2,0\}}\,.$$

In particular, $\lim_{r\to+\infty} g(r) = 0$.

Now we begin our induction scheme. We begin with f_1 which satisfies the equation (26), that is:

$$f_1'' + \frac{f_1'}{r} - n^2\frac{f_1}{r^2} + DF(f_0)f_1 = b_1(r)$$

with $b_1(r) = f_0(r)v_0^2(r)$. We want to apply Lemma 3.3 and for that we check that $\mathcal{E}[b_1] \in \mathrm{O}^3_{n-1}$. We point out that, by Proposition 3.2, $b_1 = f_0v_0^2 \in \mathrm{O}^{2,2}_{n+2}$, $b_1' \in \mathrm{O}^{3,2}_{n+1}$, $b_1'' \in \mathrm{O}^{4,2}_n$ and, consequently:

$$b_1''(r) + \frac{b_1'(r)}{r} - b_1(r)\frac{n^2}{r^2} \in \mathrm{O}^{4,2}_n\,.$$

In addition, $\big[DF(f_0(r)) + d\big]b_1(r) = \mathrm{O}\left(r^{n+2}\right)$ as $r \to 0$ and, since $DF(1) = \lambda'(1) = -d$, and using Proposition 3.2 for the asymptotics of f_0 as $r \to \infty$

$$\big[DF(f_0(r)) + d\big] = \mathrm{O}\left(f_0(r) - 1\right) = \mathrm{O}\left(r^{-2}\right) \tag{44}$$

and this gives $\big[DF(f_0(r)) + d\big]b_1(r) = \mathrm{O}\left(r^{-4}\log^2 r\right)$. Therefore we conclude that $\mathcal{E}[b_1] \in \mathrm{O}^{4,2}_n \subset \mathrm{O}^3_{n-1}$. Then, Lemma 3.3 gives the existence of a solution f_1 of problem (38) for $k = 1$ with $\Delta f_1 = f_1 + d^{-1}b_1$ satisfying

$$\Delta f_1 \in \mathrm{O}^3_n \quad \Delta f_1' \in \mathrm{O}^3_{n-1}, \quad \text{and} \quad \Delta f_1'' \in \mathrm{O}^3_{\max\{n-2,0\}},$$

which gives:

$$f_1 \in \mathrm{O}^{2,2}_n \quad f_1' \in \mathrm{O}^{3,2}_{n-1}, \quad \text{and} \quad f_1'' \in \mathrm{O}^3_{\max\{n-2,0\}}\,.$$

Now we deal with v_1 and Ω_1. As we state in (41),

$$v_1(r) = \left(rf_0^2(r)\right)^{-1}\int_0^r tf_0(t)\big(c_1(t) - f_0(t)\Omega_1\big)\,dt,$$

with c_1 defined in Proposition 3.1, formula (28). Using that $f_1 \in \mathrm{O}^{2,2}_n$, $f_1' \in \mathrm{O}^{3,2}_{n-1}$, along with $v_0 \in \mathrm{O}^{1,1}_1$ and $v_0' \in \mathrm{O}^{2,1}_0$, we have that $c_1 \in \mathrm{O}^{2,2}_n$. Therefore, v_1 will be a bounded solution if and only if $rf_0(r)\big(c_1(r) - f_0(r)\Omega_1\big)$ is a bounded function. This implies that

$$0 = \lim_{r\to+\infty} c_1(r) - f_0(r)\Omega_1 = \Omega_1.$$

Hence we actually have that

$$v_1(r) = \left(rf_0^2(r)\right)^{-1}\int_0^r tf_0(t)c_1(t)\,dt.$$

Now we need to compute the asymptotic behaviour of v_1. Clearly, for $r \to 0$, since $c_1 \in \mathrm{O}_n^{2,2}$, $v_1(r) = \mathrm{O}\,(r)$. Now we deal with $r \to +\infty$. We notice that, if r_0 is big enough,

$$\left| \int_{r_0}^{r} t f_0(t) c_1(t)\, dt \right| \le C \int_{r_0}^{r} \frac{\log^2 t}{t}\, dt \le C \log^3 r.$$

Then

$$|v_1(r)| \le C \frac{\log^3 r}{r}, \qquad \text{as } r \to +\infty.$$

Summarizing, $v_1 \in \mathrm{O}_1^{1,3}$. Moreover, from (39) with $k = 1$:

$$f_0(r) v_1'(r) = c_1(r) - 2 f_0'(r) v_1(r) - r^{-1} f_0(r) v_1(r)$$

which implies that $v_1' \in \mathrm{O}_0^{2,3}$. We can also deduce that $v_1'' \in \mathrm{O}_0^{3,3}$.

Now we state the induction hypothesis: the unique bounded solution f_{k-1}, $k \ge 2$, of problem (38) satisfies

$$f_{k-1} \in \mathrm{O}_n^{2,2(k-1)}, \;\; f_{k-1}' \in \mathrm{O}_{n-1}^{3,2(k-1)}, \;\; f_{k-1}'' \in \mathrm{O}_{\max\{n-2,0\}}^{3}\,. \tag{45}$$

Moreover, problem (39) has bounded solution v_{k-1} if and only if $\Omega_{k-1} = 0$ and in this case,

$$v_{k-1} \in \mathrm{O}_1^{1,2k-1}, \;\; v_{k-1}' \in \mathrm{O}_0^{2,2k-1}, \;\; v_{k-1}'' \in \mathrm{O}_0^{3,2k-1}\,. \tag{46}$$

We begin first by checking that $b_k \in \mathrm{O}_{n+1}^{2,2k}$ and $c_k \in \mathrm{O}_n^{2,2k}$. Indeed, by induction hypothesis (45) and (46) and formula (24) for b_k, we have that

$$b_k \in \mathrm{O}_{2n}^{2,2k} \cap \mathrm{O}_{n+2}^{2,2k} \subset \mathrm{O}_{n+1}^{2,2k}\,.$$

We emphasize that if $n = 1$, $2n < n + 2$, but if $n \ge 2$, $2n \ge n + 2$. To unify both cases we have considered $b_k \in \mathrm{O}_{n+1}^{2,2k}$. Analogously one see that $c_k \in \mathrm{O}_n^{2,2k}$.

In order to compute the orders for b_k' and c_k' we take into account that, by induction hypothesis if $l \le k-1$, the functions $f_l'(r), v_l'(r), v_l''(r)$ are of order of $f_l(r) r^{-1}$, $v_l(r) r^{-1}$ and $v_l(r) r^{-2}$ respectively, so the same happens for the products of these functions. Moreover, $f_l'' \in \mathrm{O}_{\max\{n-2,0\}}^{3}$, for $l \le k-1$. Then, tedious but easy computations yield:

$$b_k \in \mathrm{O}_{n+1}^{2,2k}, \;\; b_k' \in \mathrm{O}_n^{3,2k}, \;\; b_k'' \in \mathrm{O}_{n-1}^{3} \tag{47}$$

and

$$c_k \in \mathrm{O}_n^{2,2k}, \;\; c_k' \in \mathrm{O}_{n-1}^{3,2k}\,. \tag{48}$$

The first consequence is that $\mathcal{E}[b_k] \in \mathrm{O}_{n-1}^{3}$ and hence by Lemma 3.3 there exists a unique solution f_k of problem (38) satisfying that

$$f_k + d^{-1} b_k \in \mathrm{O}_n^{3}, \;\; f_k' + d^{-1} b_k' \in \mathrm{O}_{n-1}^{3}, \;\; f_k'' + d^{-1} b_k'' \in \mathrm{O}_{\max\{n-2,0\}}^{3}$$

and taking into account the expansions of b_k in (47), the induction hypothesis (45) is fullfilled for f_k.

Finally we deal with v_k. We proceed likewise as v_1. From identity (41),

$$v_k(r) = \big(r f_0^2(r)\big)^{-1} \int_0^r t f_0(t)\big(c_k(t) - f_0(t)\Omega_k\big)\, dt,$$

v_k will be a bounded solution if and only if $r f_0(r)\big(c_k(r) - f_0(r)\Omega_k\big)$ is a bounded function and consequently, since $c_k \in \mathrm{O}_n^{2,2k}$,

$$0 = \lim_{r\to+\infty} c_k(r) - f_0(r)\Omega_k = \Omega_k.$$

Therefore the induction hypothesis for Ω_k is also satisfied. We rewrite v_k as

$$v_k(r) = \big(r f_0^2(r)\big)^{-1} \int_0^r t f_0(t) c_k(t)\, dt,$$

and compute the asymptotic behaviour of v_k. Since $c_k(r), f_0(r) = \mathrm{O}\left(r^n\right)$ as $r \to 0$, one deduces that $v_k(r) = \mathrm{O}\left(r\right)$. As in the case $k = 1$, if r_0 is big enough,

$$\left|\int_{r_0}^r t f_0(t) c_k(t)\, dt\right| \le C \int_{r_0}^r \frac{\log^{2k} t}{t}\, dt \le C\big(\log r\big)^{2k+1}$$

and hence

$$|v_k(r)| \le C \frac{\log^{2k+1} r}{r}, \qquad \text{as } r \to +\infty.$$

Summarizing, $v_k \in \mathrm{O}_1^{1,2k+1}$. Moreover,

$$f_0(r) v_k'(r) = c_k(r) - 2 f_0'(r) v_k(r) - r^{-1} f_0(r) v_k(r)$$

implies that $v_k' \in \mathrm{O}_0^{2,2k+1}$ and we finally deduce that $v_k'' \in \mathrm{O}_0^{3,2k+1}$ by using (48).

This ends the proof of Theorem 2.2.

3.4. **Proof of Lemma 3.3.** We first write equation (42) in a more suitable way, i.e. as a fixed point equation. Adding and subtracting the term $d\,g$, where $d = -\lambda'(1)$, which is positive since $\lambda'(1) < 0$, performing the change of variables $s = \sqrt{d}\,r$ and denoting by $\tilde g(s) = g(s/\sqrt{d})$, $\tilde h(s) = d^{-1} h(s/\sqrt{d})$ yields

$$\begin{aligned} &\tilde g''(s) + \frac{\tilde g'(s)}{s} - \tilde g(s)\left(\frac{n^2}{s^2} + 1\right) = \tilde h(s) - \tilde g(s)\left[\frac{DF(\tilde f_0(s))}{d} + 1\right] \\ &\tilde g(0) = 0, \qquad \tilde g(s) \ \text{bounded} \ s \ge 0, \end{aligned} \tag{49}$$

where we call $\tilde f_0(s) = f_0(s/\sqrt{d})$.

As we showed in (44)

$$d^{-1} DF(\tilde f_0(s)) + 1 = \mathrm{O}\left(s^{-2}\right), \quad \text{as} \quad s \to +\infty. \tag{50}$$

This implies that the dominant term of equation (49) as $s \to \infty$ is the singular equation $-\tilde{g}(s) = \tilde{h}(s)$, therefore, it is natural to write $\tilde{g} = -\tilde{h} + \Delta g$ with Δg being a solution of

$$\Delta g'' + \frac{\Delta g'}{s} - \Delta g \left(\frac{n^2}{s^2} + 1 \right) = -\tilde{\mathcal{E}}[\tilde{h}](s) - \Delta g \left(\frac{DF(\tilde{f}_0(s))}{d} + 1 \right), \tag{51}$$

where $\tilde{\mathcal{E}}[\tilde{h}]$ defined by

$$\tilde{\mathcal{E}}[\tilde{h}](s) = \frac{1}{d^2}\mathcal{E}[h](s/\sqrt{d}) = \tilde{h}''(s) + \frac{\tilde{h}'(s)}{s} - \tilde{h}(s)\frac{n^2}{s^2} + \left[\frac{DF(\tilde{f}_0(s))}{d} + 1 \right] \tilde{h}(s).$$

Recall that the operator $\mathcal{E}$ is defined in the statement of the lemma. The boundary conditions are $\Delta g(0) = 0$ and $\Delta g(s)$ is bounded for $s \geq 0$.

Our goal now is to write equation (51) as a fixed point equation. We emphasize that the dominant part of this equation is the left hand side. Indeed, on the one hand, using (50), one sees that the linear term in the right hand side of equation (51), contributes a small quantity to the equation for large values of s, being the left hand side of equation (51) the dominant part as $s \to \infty$. On the other hand, as $s \to 0$, even if this linear term is of order one, the dominant part of equation (49) is provided by the first three terms of the left hand side, that is $\Delta g''(s) + \Delta g'(s)/s - \Delta g(s) n^2/s^2$, and so the right hand side in (49) is also relatively small for small values of s.

To obtain a fixed point equation we note that the homogeneous modified Bessel equation:

$$\varphi''(s) + \frac{\varphi'(s)}{s} - \varphi(s) \left(\frac{n^2}{s^2} + 1 \right) = 0$$

has two well-known linearly independent solutions, namely $I_n(s)$ and $K_n(s)$ known as the modified Bessel functions of the first and second kind respectively (see [AS64]). Hence, a fundamental matrix of solutions of the homogeneous equation corresponding to equating to zero the left hand side in (49) reads,

$$M = \begin{pmatrix} K_n(s) & I_n(s) \\ K_n'(s) & I_n'(s) \end{pmatrix},$$

whose Wronskian is known to be $W(K_n(s), I_n(s)) = 1/s$. We denote by

$$\mathcal{R}[\Delta g](s) = \tilde{\mathcal{E}}[\tilde{h}](s) + \Delta g(s) \left(\frac{DF(\tilde{f}_0(s))}{d} + 1 \right). \tag{52}$$

We recall here that Δg has to be a bounded solution of problem (51) with boundary condition $\Delta g(0) = 0$. Therefore, using the variation of parameters formula, equation (51) becomes a fixed point equation:

$$\Delta g(s) = \mathcal{F}[\Delta g](s) := K_n(s) \int_0^s \xi I_n(\xi) \mathcal{R}[\Delta g](\xi)\, d\xi + I_n(s) \int_s^\infty \xi K_n(\xi) \mathcal{R}[\Delta g](\xi)\, d\xi. \tag{53}$$

In order to prove the existence of the solution of (53) (and consequently of problem (49)), we will prove that the linear operator $\mathcal{F}$ is contractive in some appropriate Banach space $\mathcal{X}$. However to guarantee the uniqueness of this solution in the space of bounded functions, we need to carefully study the following linear operator:

$$\mathcal{T}[\psi](s) = K_n(s)\int_0^s \xi I_n(\xi)\psi(\xi)\,d\xi + I_n(s)\int_s^\infty \xi K_n(\xi)\psi(\xi)\,d\xi, \tag{54}$$

where ψ is a function defined on $J = [0, +\infty)$. We notice that $\mathcal{F} = \mathcal{T} \circ \mathcal{R}$.

The operators $\mathcal{T}$ and $\mathcal{F}$ are studied in the lemmas 3.4 and 3.5 whose proofs are deferred to the end of this section.

Lemma 3.4. *Let $\mathcal{T}$ be the linear operator defined in* (54)*. Let ψ be a function defined on $J = [0, +\infty)$. We take $0 \le m < n-1$ and $l \ge 0$. Then*

$$\psi = \mathrm{O}^l_m \Longrightarrow \mathcal{T}[\psi] = \mathrm{O}^l_{m+2},$$

where the notation O^l_m was introduced in (13)*. In particular, if ψ is bounded, then $\mathcal{T}(\psi) \in \mathrm{O}^0_2$.*

In the cases $\psi = \mathrm{O}^l_{n-1}$ or $\psi = \mathrm{O}^l_n$ we can only conclude that $\mathcal{T}[\psi] = \mathrm{O}^l_n$.

In addition, if $\psi \in \mathcal{C}^i(J)$, then $\mathcal{T}[\psi] \in \mathcal{C}^{i+1}(J)$ and

$$\psi = \mathrm{O}^l_m \Longrightarrow \mathcal{T}[\psi]' = \mathrm{O}^l_{m+1}\,.$$

In the cases $\psi = \mathrm{O}^l_{n-1}$ or $\psi = \mathrm{O}^l_n$ we conclude $\mathcal{T}[\psi]' = \mathrm{O}^l_{n-1}$.

We now define the Banach space where the solution Δg will belong. We consider the weight function

$$w(s) = f_0'\big(s/\sqrt{d}\big) \tag{55}$$

and the functional space

$$\mathcal{X} = \{\varphi : J \to \mathbb{R}, \quad \varphi \in \mathcal{C}^0(J), \quad \left|\frac{\varphi(s)}{w(s)}\right| < +\infty\}. \tag{56}$$

We endow $\mathcal{X}$ with the norm

$$\|\varphi\|_w = \sup_{s\ge 0}\left|\frac{\varphi(s)}{w(s)}\right|,$$

and it becomes a Banach space. In addition, since by Proposition 3.2, $w \in \mathrm{O}^3_{n-1}$,

$$\mathcal{X} = \mathrm{O}^3_{n-1} \cap \mathcal{C}^0(J). \tag{57}$$

Lemma 3.5. *For any given $\varphi \in \mathcal{X}$, let $\mathcal{F}_\varphi$ be the linear operator defined by* (53)*:*

$$\mathcal{F}_\varphi[g](s) = K_n(s)\int_0^s \xi I_n(\xi)\mathcal{R}_\varphi[g](\xi)\,d\xi + I_n(s)\int_s^\infty \xi K_n(\xi)\mathcal{R}_\varphi[g](\xi)\,d\xi \tag{58}$$

where, analogously to (52)*, we denote* $\mathcal{R}_\varphi$ *by*

$$\mathcal{R}_\varphi[g](s) = \varphi(s) + g(s)\left(\frac{DF(\tilde{f}_0(s)}{d} + 1\right).$$

Then,

(i) *If* $g \in \mathcal{X}$*, then* $\mathcal{F}_\varphi[g] \in \mathcal{C}^1(J)$ *and* $\mathcal{F}_\varphi[g], \mathcal{F}_\varphi[g]' \in \mathcal{X}$*. In fact* $\mathcal{F}_\varphi[g](s) \in \mathrm{O}^3_n$*.*

(ii) $\mathcal{F}_\varphi$ *is contractive in* $\mathcal{X}$*.*

End of the proof of Lemma 3.3. We have to deal with both, existence and uniqueness of solutions of problem (49). We recall that we look for $\tilde{g}$ as $\tilde{g} = -\tilde{h} + \Delta g$, with Δg being a solution of the fixed point equation $\Delta g = \mathcal{F}[\Delta g] = \mathcal{T}[\mathcal{R}[\Delta g]]$ given in (53). For the existence we will use mainly Lemma 3.5 where $\varphi(s) = \tilde{\mathcal{E}}[\tilde{h}](s)$. Then, hypothesis (43) of Lemma 3.3 and the fact that $\tilde{f}_0' \in \mathrm{O}^3_{n-1}$, assure that $\tilde{\mathcal{E}}[\tilde{h}]$ belongs to $\mathcal{X}$ and henceforth Lemma 3.5 provides us with a solution $\Delta g \in \mathcal{X}$ such that $\Delta g \in \mathrm{O}^3_n$, $\Delta g' \in \mathrm{O}^3_{n-1}$. In addition, since Δg is a solution of the differential equation (51), $\Delta g'' \in \mathrm{O}^3_{\max\{n-2,0\}}$.

Now it only remains to check that $\tilde{g} = -\tilde{h} + \Delta g$ is the unique bounded solution of our problem or equivalently, we see that Δg is the only bounded solution of (51). Let $\Delta\bar{g}$ be a bounded solution of equation (51). Then it has to be solution of the fixed point equation $\Delta\bar{g} = \mathcal{F}[\Delta\bar{g}] = (\mathcal{T} \circ \mathcal{R})[\Delta\bar{g}]$. We note that

$$\mathcal{R}[\Delta\bar{g}](s) \leq C|\Delta\bar{g}(s)| + |\tilde{\mathcal{E}}[\tilde{h}](s)|.$$

Therefore, since at least $\Delta\bar{g}$ is bounded and $\tilde{\mathcal{E}}[\tilde{h}] \in \mathcal{X}$, Lemma 3.4 with $l = m = 0$ implies that $\Delta\bar{g}(s) = \mathrm{O}\left(s^2\right)$ as $s \to 0$, then applying iteratively this lemma, we obtain that $\Delta\bar{g}(s) = \mathrm{O}\left(s^n\right)$ as $s \to 0$. In particular, since $w(s) \in \mathrm{O}^3_{n-1}$:

$$|\Delta\bar{g}(s)| \leq Cw(s), \qquad \text{as} \qquad s \to 0. \tag{59}$$

Now we study the behaviour of $\Delta\bar{g}$ as $s \to +\infty$. We first recall that, according to (50) $d^{-1}DF(\tilde{f}_0(s)) + 1 = O(s^{-2})$. Then, since $\tilde{\mathcal{E}}[\tilde{h}] \in \mathcal{X} \subset \mathrm{O}^3_{n-1}$ and $\Delta\bar{g}(s) \in \mathrm{O}^0_n$, we conclude that $\mathcal{R}[\Delta\bar{g}] \in \mathrm{O}^2_{n-1}$. Now we apply Lemma 3.4 and we obtain that $\Delta\bar{g} = \mathcal{F}[\overline{\Delta g}] = \mathcal{T}\big[\mathcal{R}[\Delta\bar{g}]\big] \in \mathrm{O}^2_n$. Therefore, repeating the previous argumentation, since $\tilde{\mathcal{E}}[\tilde{h}] \in \mathrm{O}^3_{n-1}$ and that $\Delta\bar{g} \in \mathrm{O}^2_n$ we get that $\Delta\bar{g} \in \mathrm{O}^3_n$. Hence, as $\Delta\bar{g} \in \mathcal{X}$ and $\mathcal{F}$ is a contractive operator over $\mathcal{X}$, $\Delta\bar{g} = \Delta g$. ☐

The remaining part of this Section is devoted to prove the technical lemmas 3.4 and 3.5.

3.4.1. *Proof of Lemma 3.4.* Let $m < n-1$, $l \geq 0$ and $\psi : J = [0, +\infty) \to \mathbb{R}$ be a function in O^l_m. To study the behavior of $\mathcal{T}[\psi]$ (see (54)) as $s \to 0$ and $s \to \infty$, we recall the asymptotic expansions of the modified Bessel functions K_n, I_n and their derivatives.

When $s \to 0$ one has:

$$K_n(s) \sim \frac{\Gamma(n)}{2}(s/2)^{-n} \quad I_n(s) \sim \frac{1}{\Gamma(n+1)}(s/2)^n,$$

$$K_n'(s) \sim -\frac{n\Gamma(n)}{4}(s/2)^{-n-1}, \quad I_n'(s) \sim \frac{n}{2\Gamma(n+1)}(s/2)^{n-1}.$$

And when $s \to \infty$:

$$K_n(s) \sim e^{-s}\sqrt{\pi/2s}, \quad I_n(s) \sim e^s/\sqrt{2\pi s}$$

$$K_n'(s) \sim -e^{-s}\sqrt{\pi/2s}, \quad I_n'(s) \sim e^s/\sqrt{2\pi s}.$$

From now on we will use the expansions of the Bessel functions without explicit mention.

We start by proving the behaviour of $\mathcal{T}[\psi]$ as $s \to 0$. As $\psi \in \mathrm{O}^l_m$, there exists $C > 0$ such that $|\psi(s)| \leq Cs^m$ for any $s \in J$. Let $s_0 > 0$ be such that the above expansions for $s \to 0$ are true for $0 \leq s < s_0$. We have that

$$\begin{aligned}
|\mathcal{T}[\psi](s)| \leq & CK_n(s)\int_0^s \xi I_n(\xi)\xi^m \, d\xi + CI_n(s)\int_s^{+\infty} \xi K_n(\xi)\xi^m \, d\xi \\
\leq & c\left(s^{-n}\int_0^s \xi^{n+1+m} \, d\xi + s^n \int_s^{s_0} \xi^{-n+1+m} \, d\xi + s^n \int_{s_0}^{+\infty} \xi^{1+m} K_n(\xi) \, d\xi\right) \\
\leq & \bar{c} \max\{s^{m+2}, s^n\} = \bar{c}s^{m+2}.
\end{aligned}$$

where $c, \bar{c}$ are generic constants depending only on n, s_0. We have used that, by hypothesis, $m < n-1$ and that $\int_{s_0}^{+\infty} \xi^{1+m} K_n(\xi) \, d\xi$ is bounded.

We proceed likewise with the behavior of $\mathcal{T}[\psi](s)$ as $s \to \infty$. We take $s_1 > 0$ be such that the expansions of the Bessel functions for $s \to \infty$ are true for $s > s_1$. As $\psi \in \mathrm{O}^l_m$, there exists $\overline{C}$ be such that $|\psi(s)| \leq \overline{C}s^{-l}$ for $s > s_1$ and $|\psi(s)| \leq \overline{C}$ for any $s \in J$. We obtain

$$\begin{aligned}
|\mathcal{T}[\psi](s)| \leq & \overline{C}K_n(s)\int_0^s \xi I_n(\xi)\xi^{-l} \, d\xi + \bar{C}I_n(s)\int_s^{+\infty} \xi K_n(\xi)\xi^l \, d\xi \\
\leq & ce^{-s}s^{-1/2}\left(\int_0^{s_1} \xi I_n(\xi) \, d\xi + \int_{s_1}^s \xi^{-l+1/2}e^{\xi} \, d\xi + \int_s^{+\infty} \xi^{-l+1/2}e^{-\xi} \, d\xi\right) \\
\leq & \bar{c}s^{-l},
\end{aligned}$$

where, as before, the values of $c, \bar{c}$ only depend on n, s_1. In conclusion $\mathcal{T}[\psi] \in \mathrm{O}^l_{m+2}$. In particular, applying the above inequalities for $m = l = 0$, that is ψ bounded, we have that $\mathcal{T}[\psi] \in \mathrm{O}^0_2$.

In addition, if ψ is continuous, and since every integral in the definition of $\mathcal{T}$ is uniformly convergent, we have that

$$\mathcal{T}[\psi]'(s) = K_n'(s)\int_0^s \xi I_n(\xi)\psi(\xi) \, d\xi + I_n'(s)\int_s^{\infty} \xi K_n(\xi)\psi(\xi) \, d\xi.$$

is also a continuous and bounded function. We proceed as above to check the asymptotic expansions for $\mathcal{T}[\psi]'$.

3.4.2. *Proof of Lemma 3.5.* We notice that $\mathcal{F}_\varphi = \mathcal{T} \circ \mathcal{R}_\varphi$. Let $g \in \mathcal{X}$. It is clear that $\mathcal{R}_\varphi[g] \in \mathcal{X}$ and, by (57), $g, \mathcal{R}_\varphi[g] \in \mathrm{O}^3_{n-1}$. In consequence, the first item is a straightforward consequence of Lemma 3.4.

To prove (ii) we need to show that there exists a constant $0 < K < 1$ such that, for any $g_1, g_2 \in \mathcal{X}$, $\|\mathcal{F}_\varphi[g_1] - \mathcal{F}_\varphi[g_2]\|_w \leq K\|g_1 - g_2\|_w$. We first point out that, since $0 \leq \tilde{f}_0(s) \leq 1$ and by hypothesis (A2), $\partial_x^2(x\lambda(x)) < 0$, the function $DF(x) = x\lambda'(x) + \lambda(x)$ is decreasing. Therefore, using that by hypothesis (A1) $\lambda(1) = 0$:

$$-d = \lambda'(1) + \lambda(1) = DF(1) < DF(\tilde{f}_0(s)) \leq DF(0) = \lambda(0) = 1,$$

which gives

$$0 < \frac{DF(\tilde{f}_0(s))}{d} + 1 < \frac{1}{d} + 1. \tag{60}$$

Now we find that

$$\begin{aligned} |\mathcal{F}_\varphi[g_1](s) - \mathcal{F}_\varphi[g_2](s)| \leq & K_n(s) \int_0^s \xi I_n(\xi) \left(\frac{DF(\tilde{f}_0(\xi))}{d} + 1 \right) |g_1(\xi) - g_2(\xi)| \, d\xi \\ & + I_n(s) \int_s^\infty \xi K_n(\xi) \left(\frac{DF(\tilde{f}_0(\xi))}{d} + 1 \right) |g_1(\xi) - g_2(\xi)| \, d\xi \\ & \leq \|g_1 - g_2\|_w T(s) \end{aligned}$$

where the function T is defined by

$$\begin{aligned} T(s) := K_n(s) & \int_0^s \xi I_n(\xi) \left(\frac{DF(\tilde{f}_0(\xi))}{d} + 1 \right) w(\xi) \, d\xi \\ & + I_n(s) \int_s^\infty \xi K_n(\xi) \left(\frac{DF(\tilde{f}_0(\xi))}{d} + 1 \right) w(\xi) \, d\xi. \end{aligned} \tag{61}$$

We first observe that $T(s) > 0$ $\forall s$ since both $K_n(s)$, $I_n(s)$ are positive, the weight function (see (55)) $w(s) > 0$ and inequality (60).

We now want to show that $\|T\|_w < 1$. We begin by rewriting T in a more appropriate way. Concretely, we will check that

$$\begin{aligned} T(s) &= w(s) - \mathcal{T}[h_0](s) \\ &= w(s) - K_n(s) \int_0^s \xi I_n(\xi) h_0(\xi) \, d\xi - I_n(s) \int_s^{+\infty} \xi K_n(\xi) h_0(\xi) \, d\xi \end{aligned} \tag{62}$$

being $\mathcal{T}$ the linear operator defined in Lemma 3.4 and

$$h_0(s) = \frac{\sqrt{d}}{s^3} \left[2n^2 \tilde{f}_0(s) - \frac{s}{\sqrt{d}} f_0' \left(\frac{s}{\sqrt{d}} \right) \right]. \tag{63}$$

To prove expression (62) we deal with the differential equation that $f_0'(r)$ satisfies. Indeed, since $f_0(r)$ is a solution of equation (20), $f_0'(r)$ is a solution of the nonhomogeneous linear equation:

$$\varphi''(r) + \frac{\varphi'(r)}{r} - \varphi(r)\frac{n^2}{r^2} + DF(f_0(r))\varphi(r) = -\frac{2n^2}{r^3}f_0(r) + \frac{1}{r^2}f_0'(r). \tag{64}$$

Performing the change $\psi(s) = \varphi(s/\sqrt{d})$ to this equation and taking into account that $\tilde{f}_0(s) = f_0(s/\sqrt{d})$, we get that $w(s) = f_0'(s/\sqrt{d})$ is a solution of

$$\psi''(s) + \frac{\psi'(s)}{s} - \psi(s)\frac{n^2}{s^2} + \frac{DF(\tilde{f}_0(s))}{d}\psi(s) = -\frac{2n^2\sqrt{d}}{s^3}\tilde{f}_0(s) + \frac{1}{s^2}f_0'(s/\sqrt{d}). \tag{65}$$

We define

$$\mathcal{L}[\psi](s) = \psi''(s) + \frac{\psi'(t)}{s} - \psi(s)\left(\frac{n^2}{s^2} + 1\right). \tag{66}$$

We notice that $\mathcal{L}[K_n] = \mathcal{L}[I_n] = 0$ and that equation (65), for w, can be rewritten as

$$\left(\frac{DF(\tilde{f}_0(s))}{d} + 1\right) w(s) = -\mathcal{L}[w](s) - h_0(s). \tag{67}$$

The linear differential operator $\mathcal{L}$ satisfies that, upon integrating by parts,

$$-\int_a^b \xi B_n(\xi)\mathcal{L}[\psi](\xi)\, d\xi = -\, \xi\psi'(\xi)B_n(\xi)\big|_a^b + \xi\psi(\xi)B_n'(\xi)\big|_a^b, \tag{68}$$

being either $B_n = K_n$ or $B_n = I_n$. This property was strongly used in [AB11]. Using that w satisfies equation (67), property (68) and that $s(I_n'(s)K_n(s) - K_n'(s)I_n(s)) = 1$, we have that definition (61) of T becomes

$$\begin{aligned} T(s) =& -K_n(s)\int_0^s \xi I_n(\xi)\mathcal{L}[w](\xi)\, d\xi - I_n(s)\int_s^\infty \xi K_n(\xi)\mathcal{L}[w](\xi)\, d\xi. \\ & - K_n(s)\int_0^s \xi I_n(\xi)h_0(\xi)\, d\xi - I_n(s)\int_s^{+\infty} \xi K_n(\xi)h_0(\xi)\, d\xi \\ & = w(s) - \mathcal{T}[h_0](s) \end{aligned}$$

and (62) is proven.

Since by Proposition 3.2, for any $r \geq 0$, $rf_0'(r) \leq n^2 f_0(r)$, we have that h_0, defined in (63), satisfies that $h_0(s) > 0$ if $s > 0$, and therefore $\mathcal{T}[h_0](s) > 0$, $s > 0$. Consequently:

$$0 \leq T(s) < w(s) \quad s > 0.$$

Now, in order to check that $\|T\|_w < 1$ it only remains to see that

$$\lim_{s\to 0}\frac{\mathcal{T}[h_0](s)}{w(s)} \neq 0, \qquad \lim_{s\to+\infty}\frac{\mathcal{T}[h_0](s)}{w(s)} \neq 0.$$

Indeed, recalling again definition (55) of w, and using Proposition 3.2, we have that

$$\lim_{s\to 0} w(s)s^{-n+1} = n\alpha d^{-(n-1)/2}, \quad \lim_{s\to+\infty} s^3 w(s) = 2n^2\sqrt{d},$$

$$\lim_{s\to 0} h_0(s)s^{-n+3} = \frac{n(2n-1)\alpha}{d^{(n-1)/2}}, \quad \lim_{s\to+\infty} s^3 h_0(s) = 2n^2\sqrt{d}.$$

Let $s_0 >$ be small enough. By applying Hôpital's rule

$$\begin{aligned}
\lim_{s\to 0} \frac{\mathcal{T}[h_0](s)}{w(s)} &= \frac{d^{(n-1)/2}}{2n^2\alpha} \lim_{s\to 0} \left(s^{-2n+1} \int_0^s \xi^{n+1} h_0(\xi) + s \int_s^{s_0} \xi^{-n+1} h_0(\xi) \right) \\
&= \frac{d^{(n-1)/2}}{2n^2\alpha} \lim_{s\to 0} \left(\frac{s^{n+1} h_0(s)}{(2n-1)s^{2n-2}} + s^{-n+3} h_0(s) \right) \\
&= \frac{1}{2n} + \frac{2n-1}{2n} = 1.
\end{aligned}$$

Now we deal with $s \to +\infty$. Let then $s_0 > 0$ be big enough. Then,

$$\begin{aligned}
\lim_{s\to+\infty} \frac{\mathcal{T}[h_0](s)}{w(s)} &= \frac{1}{4n^2\sqrt{d}} \lim_{s\to+\infty} \left(s^{5/2} e^{-s} \int_{s_0}^s e^{\xi} \xi^{1/2} h_0(\xi)\, d\xi + s^{5/2} e^{s} \int_s^{+\infty} e^{-\xi} \xi^{1/2} h_0(\xi)\, d\xi \right) \\
&= \frac{1}{4n^2\sqrt{d}} \lim_{s\to+\infty} 2s^3 h_0(s) = 1.
\end{aligned}$$

Acknowledgements M. Aguareles has been supported in part by grants from the Spanish Government MTM2011-27739-C04-03, MTM2014-52402-C3-3-P and and is a member of the Catalan research group 2014SGR1083. I. Baldomá and T.M-Seara have been partially supported by the Spanish MINECO-FEDER Grant MTM2012-31714 and the Catalan Grant 2014SGR504 . T. M-Seara has been partially supported by the Russian Scientific Foundation grant 14-41-00044 and Marie Curie Action FP7-PEOPLE-2012-IRSES, BREUDS.

REFERENCES

[AB11] M. Aguareles and I. Baldomá. Structure and Gevrey asymptotic of solutions representing topological defects to some partial differential equations. *Nonlinearity*, 24(10):2813–2847, 2011.

[ACW10] M. Aguareles, S.J. Chapman, and T. Witelski. Motion of spiral waves in the complex ginzburglandau equation. *Physica D: Nonlinear Phenomena*, 239(7):348 – 365, 2010.

[AS64] M. Abramowitz and I.A. Stegun. *Handbook of Mathematical Functions*. Dover, New York, fifth edition, 1964.

[BHO97] Tomas Bohr, Greg Huber, and Edward Ott. The structure of spiral-domain patterns and shocks in the 2D complex Ginzburg-Landau equation. *Phys. D*, 106(1-2):95–112, 1997.

[GB05] Marcus R Garvie and James F Blowey. A reaction-diffusion system of lambda-omega type part ii: Numerical analysis. *European Journal of Applied Mathematics*, 16(5):621, 2005.

[Gre81] J.M. Greenberg. Spiral waves for $\lambda - \omega$ systems. *Add. Appl. Math.*, 2, 1981.

[Hag82] Patrick S. Hagan. Spiral waves in reaction-diffusion equations. *SIAM J. Appl. Math.*, 42(4):762–786, 1982.

[Kur84] Y. Kuramoto. *Chemical oscillations, waves and turbulence*, volume 19 of *Springer Series in Synergetics*. Springer-Verlag, Berlin, 1984.

[LJD91] Peralta E. Leichleiter J., Girard S. and Clapham D. Spiral calcium wave propagation and annihilation in Xenopus-Laevis Oocytes. *Science*, 252(5002):123–126, April 5 1991.

[NK81] L.N. Howard N. Kopell. Target pattern and spiral solutions to reaction-diffusion equations with more than one space dimension. *Add. Appl. Math.*, 2, 1981.

[Sch98] Arnd Scheel. Bifurcation to spiral waves in reaction-diffusion systems. *SIAM J. Math. Anal.*, 29(6):1399–1418 (electronic), 1998.

[Win72] A.T. Winfree. Spiral Waves of Chemical Activity. *Science*, 175(4022):634–&, 1972.

[ZZ70] A.N. Zaikin and A.M. Zhabotinsky. Concentration wave propagation in two-dimensional liquid phase self-oscillating system. *Nature*, 225:535–537, 1970.